

1-4252k

MYOCARDIAL GRAFTS AND CELLULAR  
COMPOSITIONS USEFUL FOR SAME

BACKGROUND OF THE INVENTION

The present invention resides generally in the field of cardiology, and more particularly relates to stable myocardial grafts and methods and cellular compositions useful for achieving such grafts.

As further background, organ transplantation has been widely used to replace diseased, nonfunctional tissue. More recently, cellular transplantation to augment deficiencies in host tissue function has emerged as a potential therapeutic paradigm. One example of this approach is the well publicized use of fetal tissue in individuals with Parkinsonism (reviewed in (1), see reference list, infra.), where dopamine secretion from transplanted cells alleviates the deficiency in patients. In other studies, transplanted myoblasts from-uneffected siblings fused with endogenous myotubes in Duchenne's patients; importantly the grafted myotubes expressed wild-type dystrophin (2).

"Express Mail" label number TR 718 436 378 US  
Date of Deposit June 7, 1995

I hereby certify that this paper or fee is being deposited with the United States Postal Service "Express Mail Post Office to Addressee" service under 37 CFR §1.10 on the date indicated above and is addressed to the Commissioner of Patents and Trademarks, Washington, D.C. 20231.

David J. Kitterburg  
Signature of person mailing paper or fee

Despite their relevance in other areas, these earlier studies do not describe any cellular transplantation technology which can be successfully applied to the heart, where the ability to replace damaged myocardium would have obvious clinical relevance. Additionally, the use of intra-cardiac grafts to target the long-term expression of angiogenic factors and inotropic peptides would be of therapeutic value for individuals with myocardial ischemia or congestive heart failure, respectively.

In light of this background there is a need for the development of cellular transplantation technology in the heart. Desirably, such technology would not only provide stable grafts in the heart but also enable the delivery of useful recombinant proteins or other molecules directly to the heart. The present invention addresses these needs.

093701-0600  
F00000 F00000

# SUMMARY OF THE INVENTION

The applicant has established cellular grafts in the myocardium which are viable long-term. Cardiomyocytes and skeletal myoblasts have been grafted directly into the myocardium of syngeneic animals. Viable grafts were detected at least one-half year post-implantation (the latest time point assayed). The presence of the grafts was not accompanied by overt cardiac arrhythmia, and the majority of the grafts were juxtaposed directly to the host myocardium and not encapsulated. It has thus been discovered that the myocardium can serve as a stable platform for cellular transplants. These transplants can be used for the local delivery of recombinant molecules to the heart and/or for replacing diseased tissue to supplement myocardial function.

Accordingly, one preferred embodiment of the invention provides a myocardial graft in an animal which includes a stable graft of skeletal myoblasts or cardiomyocytes incorporated in myocardial tissue of the animal.

Another preferred embodiment of the invention provides a method for forming a stable myocardial graft in an animal. The inventive method includes the step of introducing skeletal myoblasts or cardiomyocytes in myocardial tissue of the animal so as to form a stable myocardial graft. The cells can be conveniently introduced, for example, by injection.

Another preferred embodiment of the invention provides a method for delivering a recombinant molecule to myocardial tissue of an animal. This method includes the step of establishing a stable graft of skeletal myoblasts or cardiomyocytes incorporated in myocardial tissue of the

animal, wherein the myoblasts or cardiomyocytes deliver the recombinant molecule to the myocardial tissue. In this embodiment the myoblasts or cardiomyocytes will carry transgenes encoding the recombinant molecule.

Another preferred embodiment of the invention provides a cellular composition comprising a substantially homogeneous population of non-immortalized cardiomyocytes. This and other cell populations can be obtained utilizing a preferred inventive method that includes (i) transfecting embryonic stem cells to introduce a marker gene enabling selection of one cell lineage from other cell lineages resulting from differentiation of the stem cells, (iii) causing the stem cells to differentiate, and (iv) selecting said one cell lineage based on the marker gene. The cells used in and resulting from such methods also form a part of the present invention.

Still another preferred embodiment of the invention provides a non-human animal having a stable graft of skeletal myoblasts or cardiomyocytes incorporated in myocardial tissue of the animal.

The invention thus provides myocardial grafts, methods and cellular compositions useful for forming myocardial grafts, and animals which have the myocardial grafts. The grafts will find use both as a vehicle for delivering therapeutic substances such as recombinant proteins and other molecules, and as a means for replacing diseased tissue to supplement myocardial function. Cellular compositions of the invention can be used directly to prepare grafts, and will also be useful in screening drug substance effects on cardiomyocytes and for expressing and obtaining recombinant proteins. Grafted animals can be used, for example, to screen the effects of

recombinant molecules on the heart.

These and other objects and advantages of the invention will be be apparent from the following description.

BRIEF DESCRIPTION OF THE FIGURE

Figure 1 is a schematic diagram illustrating DNA used to generate MHC-nLAC transgenic mice in Example 3, infra.

0997001-0600  
T00000 T00000

## DESCRIPTION OF THE PREFERRED EMBODIMENT

For the purpose of promoting an understanding of the principles of the invention, reference will now be made to certain embodiments thereof and specific language will be used to describe the same. It will nevertheless be understood that no limitation of the scope of the invention is thereby intended, such alterations, further modifications and applications of the principles of the invention as illustrated herein being contemplated as would normally occur to one skilled in the art to which the invention relates.

As indicated above, the present invention provides stable myocardial grafts of skeletal myoblasts and/or cardiomyocytes. In this regard, as used herein the term "stable myocardial graft" is intended to mean a myocardial graft whose cells are viable for a period of at least about 2 weeks. Surprisingly, such stable grafts have been readily achieved in accordance with the invention, with preferred grafts having cells viable for six months or more. Myocardial grafts of the invention can thus provide for long-term delivery of recombinant proteins or other molecules to the heart and/or long-term supplementation of myocardial tissue.

The skeletal myoblasts and cardiomyocytes used in the invention can be obtained or isolated from any suitable source. Skeletal myoblasts, including for example C2C12 skeletal myoblasts, are available from public depositories such as the American Type Culture Collection (ATCC) (Rockville, Maryland). Skeletal myoblasts can also be isolated from skeletal muscle using techniques well known to the art and literature. Cardiomyocytes useful for the invention can be obtained using techniques described in the literature (3) or using methods described more

particularly in the Examples below. Briefly, one such method involves digestion of heart tissues to obtain cardiomyocytes.

Another method involves using an appropriate marker to select specific cell lineages, such as cardiomyocytes, from other cell lineages resulting from the differentiation of embryonic stem cells (totipotent cell lines derived from the inner cell mass of blastocysts as described in (22)). The preferred method involves a positive selection scheme. Thus, a marker gene, such as a gene conferring antibiotic resistance (e.g. neomycin or hygromycin), is introduced into the stem cells under appropriate control such that expression of the gene occurs only in the desired cell lineage. For example, the marker gene can be under the control of a promoter which is active only in the desired cell lineage. Upon differentiation of the stem cells, the desired lineage is then selected based upon the marker, e.g. by contacting the mixed differentiated cells with the appropriate antibiotic to which the desired lineage has been conferred resistance. Cell lines other than the desired line will thus be killed, and a substantially pure, homogeneous population of the desired line can be recovered. In more preferred methods, two markers are introduced into the parent stem cells, one allowing selection of transfected stem cells from non-transfected cells, and one allowing selection of the desired cell lineage from other lineages. A double positive selection scheme can thus be used where each selectable marker confers antibiotic resistance. Using this selection methodology, populations comprised about 90% and even about 95-100% of the desired cell lineage can be obtained, as demonstrated in the Examples below.

To obtain grafts of the invention, the skeletal myoblasts or cardiomyocytes will be introduced into the



myocardial tissue of a living animal such as a mammal. The cells can be introduced in any suitable manner, but it is preferred that the mode of introduction be as non-invasive as possible. Thus, delivery of the cells by injection, catheterization or similar means will be more desired.

The resulting graft-bearing animals have exhibited normal sinus rhythms, indicating that the graft per se, as well as the graft-host myocardium border zone, does not induce arrhythmias. This is in stark contrast to the remodeling that frequently occurs following infarcts in humans; the border zone of the infarct may give rise to circus loops which result in clinically significant arrhythmias (4, 5).

Grafts of the invention can be proliferative or non-proliferative. For example, the AT-1 grafts established in the specific Examples below are proliferative. On the other hand, the skeletal myoblast-derived grafts formed in the Examples are non-proliferative, with the absence of tritiated thymidine uptake demonstrating that the formation of stable intra-cardiac grafts was not dependent upon sustained cell proliferation.

Preferred grafts will be characterized by the presence of direct intracellular coupling and the formation of gap junctions between host and grafted cells. Moreover, such grafts will not cause immune response in the host, and will exhibit terminal differentiation of grafted cells and a non-tumorigenic nature.

Grafts of the invention are useful inter alia to deliver therapeutic proteins and the like via secretion

from grafted cells, and to replace diseased or damaged tissue to supplement myocardial function. As examples of therapeutic protein deliveries, grafts may express angiogenic factors (as exemplified by basic and acidic Fibroblast Growth Factor; Transforming Growth Factor-Beta, Vascular Endothelial Growth Factor and Hepatocyte Growth Factor) to induce neovascularization. Similarly, grafts expressing neurotrophic agents near an infarcted region may be used to ameliorate the arrhythmogenesis associated with the border zone. These and many other candidate substances for targeted delivery to the heart will be apparent to those skilled in the area.

To promote a further understanding of the invention and its principles and advantages, the following specific Examples are provided. It will be understood that these Examples are illustrative, and not limiting, in nature.

#### EXAMPLE 1

##### Generation of Stable AT-1 Cardiomyocyte Grafts

###### A. METHODS

AT-1 Cell Culture and Myocardial Grafting Protocol. AT-1 cardiomyocytes were isolated from subcutaneous tumors by sequential collagenase digestion and cultured in PC-1 medium (Ventrex, Coon Rapids MN) containing 10% fetal calf serum as previously described in (6). Cells were labeled with 10  $\mu$ M 8-chloromethyl-4,4-difluoro-1,3,5,7,-tetramethyl-4-bora-3a,4a-diazaindecene (BODIPY, Molecular Probes, Eugene OR) for 30 min at 37°C to facilitate localization of the injection site. Immediately before injection, cells were harvested with trypsin and collagenase, washed three times with serum-free PC-1 medium and directly injected into the ventricular myocardium of syngeneic B6D2/F1 mice (Jackson

Laboratories, Bar Harbor MA) under open heart surgery as described in (7). Cells ( $4-10 \times 10^4$ ) were injected in a volume of 2-3  $\mu$ l using a plastic syringe fitted with a 30 gauge needle.

Histology. Hearts were removed following cervical dislocation and cryoprotected in 30% sucrose, embedded and sectioned at 10  $\mu$ m with a cryomicrotome as described (8). For hematoxylin and eosin (H and E) staining, sections were post fixed in acetone:methanol (1:1) and stained according to manufacturer's specifications (Sigma Diagnostics, St. Louis MO). For immuno-histology, unfixed sections were reacted with polyclonal rabbit anti-T-Ag antibodies (either 161-T, see (3) or 162-T) followed by horseradish peroxidase-conjugated goat anti-rabbit antisera (Boehringer Mannheim, Indianapolis IN), and visualized by diaminobenzidine reaction with nickel enhancement as described in (9). Monoclonal antibodies against the common leukocyte antigen (CD45; antibody M1/9.3HL, Boehringer Mannheim) and against the macrophage Mac-1 antigen (CD11b; antibody M1/70HL, Boehringer Mannheim) were used to monitor intra-cardiac graft rejection. The Mac-1 antibody has 75-90% cross reactivity with lymphocytes. After treatment with primary antibody, sections were incubated with horseradish peroxidase-conjugated rabbit anti-rat antisera (Boehringer Mannheim), and visualized by diaminobenzidine reaction with nickel enhancement. For [ $^3$ H]-thymidine incorporation, mice were given a single bolus injection of isotope (400  $\mu$ Ci at 28 Ci/mM, Amersham, Arlington Heights IL) and eighteen hours later sacrificed by cervical dislocation. The heart was removed, cryoprotected in 30% sucrose, embedded and sectioned with a cryomicrotome. Sections were post-fixed in methanol:acetone (1:1), stained with H and E, and a thin layer of photographic emulsion (Ilford L.4, Polysciences,

Warrington PA) diluted 1:1 with distilled water was applied. Sections were exposed for 5-7 days at 4°C, and developed in Kodak D-19 at 20°C for 4 minutes, washed with distilled water for 1 minute, fixed in 30% sodium thiosulfate for 10 minutes, and washed in distilled water.

Electron Microscopy (EM). Tissue blocks were fixed in 2% glutaraldehyde in 0.1M cacodylate buffer (pH 7.4) and post-fixed in 2% osmium tetroxide (Stevens Metallurgical Corp., New York NY). All other EM chemicals were obtained from Ladd Research Industries, Inc. (Burlington VT). Tissue was stained en bloc with 2% uranyl acetate in pH 5.2 maleate buffer (0.05 M), dehydrated, and embedded in Ladd LX-112. Grafts were located using 1µm sections stained with toluidine blue. After trimming, the block was thin sectioned, and stained with uranyl acetate and lead citrate. Specimens were viewed on a Phillips 400 transmission electron microscope.

Electrocardiogram (ECG) Analyses. For surface ECG records, mice were anesthetized (2.5% Avertin, 0.015 ml/g body weight, IP, Fluka Chemicals, Lake Ronkonkoma NY), surface electrodes were placed in the standard lead I position, and ECGs were recorded with a Narco Biosystems (Houston TX) high gain amplifier coupled to an A/D converter (Coulbourn Instruments, Lehigh Valley PA).

Plasma Enzyme Assay (PEA). For lactate dehydrogenase (LDH) isoform assay, plasma was isolated by retro-orbital sinus bleeds under anesthesia (2.5% Avertin, 0.015 ml/g body weight, intraperitoneally (IP)). Plasma was fractionated on 1% agarose gels (CK Isoenzyme electrophoresis system, CIBA-Corning Diagnostics, Corning N.Y.) and the LDH isoforms visualized by a TNBT-Formazan histochemical assay (LDH Assay Kit, Sigma Diagnostics, St. Louis MO).

## B. RESULTS

In these studies, AT-1 cardiomyocytes (derived from transgenic animals that expressed the T-Ag oncoprotein in the heart) were injected directly into the myocardium of syngeneic mice and the viability of the grafted material was assessed. To facilitate localization of the injection site in preliminary experiments, AT-1 cardiocytes were incubated briefly with BODIPY prior to grafting. BODIPY is a nontoxic glutathione reactive dye which permits fluorescent tracking of living cells. The graft site was easily visualized by fluorescence microscopy using a FITC cube. Subsequent experiments did not utilize BODIPY.

Fifty percent (14/28) of the animals receiving AT-1 cardiomyocyte injections developed intra-cardiac grafts. In most instances, the grafts were neither encapsulated nor surrounded by scarred myocardium. At the level of light microscopy, grafted AT-1 cardiomyocytes were observed directly juxtaposed with host cardiomyocytes. The identity of the AT-1 cardiomyocytes was confirmed by immuno-peroxidase assay using an anti-T-Ag antibody primary antibody (162-T) followed by a horseradish peroxidase conjugated secondary antibody. Specificity of the anti-T-Ag antibody has been established previously (10, 11). Black precipitate was observed over cardiomyocyte nuclei in the graft but not in the host myocardium, confirming that the graft was comprised of AT-1 cardiomyocytes. Similar results were obtained with other anti-T-Ag antibodies, and no signal was observed in the absence of primary antibody.

Viable AT-1 cardiomyocytes were observed at least as long as four months post-implantation. During this period, some degree of graft proliferation occurred;

Immunohistologic experiments were employed to determine if the intra- cardiac grafts were subject to chronic rejection. Grafts older than one month failed to react with antibodies specific for mouse leukocytes; signals observed in blood vessels located on the same section provided a positive control for the experiment. Similarly, an antibody which detects mouse macrophages and lymphocytes did not react with the intra-cardiac graft; once again positive signal was observed in a blood vessel located on the same section. Collectively, these results indicate the absence of chronic graft rejection by the syngeneic hosts. This result is supported by the observation that cyclosporine treatment (50 mg/kg body weight, administered intraperitoneally daily) did not influence significantly the frequency of intra-cardiac grafting (50% success rate, n=6). Sex of the host animal also did not appear to influence significantly the rate of graft formation (46% success rate in males, n=13; 53% success rate in females, n=15). The frequency of grafting was similar in animals examined at early time points (1-40 days post-grafting, 47%, n=15) as compared to those examined at later time points (40-120 days post-grafting, 54%, n=13). Finally, similar frequencies of intra-cardiac grafting were observed when cells were delivered to either the left ventricular free wall or the apex of the heart.

Electron microscopic analysis of the AT-1 cardiomyocyte grafts confirmed the absence of encapsulation. High power views revealed well-developed junctional complexes between adjacent cells within the graft. Graft cardiomyocytes contained numerous polyribosomes and the dedifferentiated myofibrillar ultrastructure typical of AT-1 tumors in vivo (6). Electron-dense secretory granules were also observed in the AT-1 cardiomyocyte grafts, as would be expected for myocytes of atrial origin. Host cardiomyocytes bordering the grafts had normal ultrastructure with well-formed sarcomeres. Although only a thin basement membrane separated AT-1 and host cardiomyocytes, no junctional connections between these two cell types were observed.

Surface electrocardiograms were performed to determine if the presence of AT-1 cardiomyocyte grafts influenced the autonomic rhythm. No appreciable differences were observed between records from sham animals and those which harbored grafts. In each case, the experimental animals exhibited normal sinus rhythm, with an anesthetized heart rate of approximately 400 beats per minute. Normal P-QRS coupling was maintained, indicating that the grafted AT-1 cardiomyocytes did not act as an ectopic pacemaker. This latter result is important in light of the observation that AT-1 cardiomyocytes exhibit spontaneous electrical activity both in vivo (12) and in culture (3). The absence of overt arrhythmia also indicated that graft-induced myocardial remodeling was not associated with the generation of significant circus rhythms.

In addition to surface ECG, plasma LDH levels were assessed in mice carrying AT-1 cardiomyocyte grafts. The presence of the cardiac LDH isoform in the circulation is a well established hallmark of myocardial infarction. No

cardiac LDH (isoform-1) was apparent in mouse plasma prior to grafting. After the introduction of AT-1 cardiomyocytes, there was a transient appearance of the cardiac isoform in the plasma, which most likely reflected damage to the host myocardium as well as damaged AT-1 cardiomyocytes. A transient increase in plasma skeletal LDH isoform was also observed following grafting surgery, presumably reflecting damage caused by the trans-thoracic incision. The plasma LDH profiles returned to normal by 7 days post-implantation. Thereafter, the plasma LDH profiles remained normal despite the presence of grafts.

#### EXAMPLE 2

#### Generation of Stable C2C12 Myoblast Grafts

##### A. METHODS

##### C2C12 Cell Culture and Myocardial Grafting Protocol.

C2C12 myoblasts were obtained from ATCC. Cells were maintained in the undifferentiated state by culturing at low density in high glucose Dulbecco's Modified Eagle Media (DMEM) supplemented with 20% fetal bovine serum, 1% chicken embryo extract, 100 units/ml penicillin and 100 µg/ml streptomycin. For some studies, myogenic differentiation was induced by culturing in DMEM supplemented with 2% horse serum and antibiotics. Immediately before injection, myoblasts were harvested with trypsin, washed three times with serum free DMEM and directly injected into the ventricular myocardium of adult syngeneic C3Heb/FeJ mice (Jackson Laboratories) under open heart surgery as described in (7). Cells ( $4-10 \times 10^4$ ) were injected in a volume of 2-3 µl using a plastic syringe fitted with a 30 gauge needle.

Histology. Hearts were removed, cryoprotected, embedded and sectioned as in Example 1. H and E staining and

2025 RELEASE UNDER E.O. 14176



[<sup>3</sup>H]-thymidine incorporation assays were also conducted as in Example 1. For immunohistology, methanol fixed sections (-20°C, 10 min.) were reacted with the monoclonal anti-skeletal myosin heavy chain antibody (MY-32, Sigma Chemical Corp.) followed by rhodamine-conjugated sheep anti-mouse IgG F(ab')<sub>2</sub> fragment (Boehringer Mannheim), and visualized by epifluorescence.

Electron Microscopy. EM was performed as in Example 1.

Electrocardiogram Analyses. ECG analyses were performed as in Example 1.

Plasma Enzyme Assay. PEA was performed as in Example 1.

## B. RESULTS

Several myoblast cell lines are known which, as exemplified by C2C12 cells, have the capacity to differentiate into myotubes in culture (13). C2C12 myoblasts were derived from cultured explants of injured thigh muscle of C3H mice. When maintained in serum-rich media, the myoblasts proliferate rapidly and retain an undifferentiated phenotype. However, when cultured in serum-poor media myogenic differentiation is induced. The C2C12 cells withdraw from the cell cycle and fuse, thereby forming multinucleated myotubes. Myogenic differentiation is also induced, as evidenced by the appearance of numerous muscle-specific gene products. Thus, in this model proliferation and myogenic differentiation are mutually exclusive (14). Myoblast differentiation in vitro is thought to mimic satellite cell mediated myofiber regeneration in vivo.

Myoblasts were injected directly into the myocardium of syngeneic C3Heb/FeJ mice and the viability of the

grafted material was assessed. One hundred percent (13/13) of the mice receiving intra-cardiac implants of C2C12 myoblasts developed grafts in the heart. Viable grafts were observed as long as six months post-implantation (this was the last time point assayed). In all instances, the grafted material was not encapsulated. The differentiated status of the grafted C2C12 cells was determined by immunohistological assay with an anti-myosin heavy chain antibody (MY-32). This antibody does not react with myoblasts nor with cardiac myosin heavy chain. Although differentiated C2C12 cells were observed in every heart receiving myoblast injections, the grafting efficiency of individual cells was not determined. As an additional control, hearts bearing AT-1 intra-cardiac grafts (see Example 1) were examined with the MY-32 antibody. No staining was observed, thereby ruling out the possibility that the signal seen in the C2C12 grafts was due to skeletal myosin heavy chain induction in host cardiomyocytes.

Example 1 above demonstrates that AT-1 cardiomyocytes form stable grafts in syngeneic myocardium. However, the observation that these cells retained the capacity for proliferation in vivo raised the possibility that sustained cell division might be required for successful intra-cardiac grafting. The proliferative status of the C2C12 grafts was therefore examined. Virtually no DNA synthesis (as assessed by tritiated thymidine incorporation) was observed, indicating that the majority of the grafted C2C12 cells had indeed withdrawn from the cell cycle. Examination of serial sections indicated that less than 0.1% of the cells in or near the grafts were synthesizing DNA. This result most likely reflects fibroblast proliferation during the remodeling process. As with the AT-1 grafts, immunohistological analyses of C2C12 grafts failed to detect macrophage,

inflammatory leukocyte or lymphocyte infiltration at two months post-implantation, indicating the absence of chronic graft rejection by the syngeneic hosts.

At the level of light microscopy, the C2C12 intra-cardiac grafts exhibited cellular heterogeneity with both H and E and MY-32 immunofluorescence staining. Electron microscopic analyses were employed in an effort to further characterize the cellular make-up of the C2C12 grafts. Toluidine-stained 1  $\mu$ m sections were surveyed at 100  $\mu$ m intervals to locate graft sites for EM analysis. Once localized, thin sections were prepared from the block. Cells with morphology typical of skeletal myocytes were observed throughout the graft. Abundant mitochondria localized between well developed sarcomeres were readily detected. Prominent Z bands and thick and thin filaments were observed. Occasionally, expanded t-tubules and ruffled cell membranes were detected in the grafted myocytes. In addition to well developed myocytes, a second less differentiated cell type was observed in C2C12 grafts. Most notably, these cells exhibited a large nucleus to cytoplasm ratio, with a prominent band of heterochromatin at the nuclear periphery. Moderate amounts of centrally located heterochromatin were also detected. Limited rough endoplasmic reticulum and few mitochondria were observed in these cells. Similar ultrastructural characteristics have been ascribed to satellite cells in vivo and in culture (15, 16).

Two studies were initiated to assess any deleterious effects of C2C12 intra-cardiac grafts on host heart function. In the first study, surface electrocardiograms failed to detect any appreciable differences between records from control and experimental mice. All animals examined had normal P-QRS coupling, and exhibited normal sinus rhythm with an anesthetized heart rate of

1. $\alpha$ = 0.05 (one-tailed)	
df	$\alpha$
1	0.0500
2	0.0250
3	0.0125
4	0.0075
5	0.0050
6	0.0038
7	0.0030
8	0.0025
9	0.0020
10	0.0018
11	0.0015
12	0.0013
13	0.0011
14	0.0010
15	0.0009
16	0.0008
17	0.0007
18	0.0006
19	0.0005
20	0.0005
21	0.0004
22	0.0004
23	0.0003
24	0.0003
25	0.0003
26	0.0002
27	0.0002
28	0.0002
29	0.0002
30	0.0002
31	0.0001
32	0.0001
33	0.0001
34	0.0001
35	0.0001
36	0.0001
37	0.0001
38	0.0001
39	0.0001
40	0.0001
41	0.0001
42	0.0001
43	0.0001
44	0.0001
45	0.0001
46	0.0001
47	0.0001
48	0.0001
49	0.0001
50	0.0001
51	0.0001
52	0.0001
53	0.0001
54	0.0001
55	0.0001
56	0.0001
57	0.0001
58	0.0001
59	0.0001
60	0.0001
61	0.0001
62	0.0001
63	0.0001
64	0.0001
65	0.0001
66	0.0001
67	0.0001
68	0.0001
69	0.0001
70	0.0001
71	0.0001
72	0.0001
73	0.0001
74	0.0001
75	0.0001
76	0.0001
77	0.0001
78	0.0001
79	0.0001
80	0.0001
81	0.0001
82	0.0001
83	0.0001
84	0.0001
85	0.0001
86	0.0001
87	0.0001
88	0.0001
89	0.0001
90	0.0001
91	0.0001
92	0.0001
93	0.0001
94	0.0001
95	0.0001
96	0.0001
97	0.0001
98	0.0001
99	0.0001
100	0.0001
101	0.0001
102	0.0001
103	0.0001
104	0.0001
105	0.0001
106	0.0001
107	0.0001
108	0.0001
109	0.0001
110	0.0001
111	0.0001
112	0.0001
113	0.0001
114	0.0001
115	0.0001
116	0.0001
117	0.0001
118	0.0001
119	0.0001
120	0.0001
121	0.0001
122	0.0001
123	0.0001
124	0.0001
125	0.0001
126	0.0001
127	0.0001
128	0.0001
129	0.0001
130	0.0001
131	0.0001
132	0.0001
133	0.0001
134	0.0001
135	0.0001
136	0.0001
137	0.0001
138	0.0001
139	0.0001
140	0.0001
141	0.0001
142	0.0001
143	0.0001
144	0.0001
145	0.0001
146	0.0001
147	0.0001
148	

## Generation of Stable Fetal Cardiomyocyte Grafts

Cardiomyocyte Cell Culture and Myocardial Grafting Protocol. Transgenic mice were generated which carry a fusion gene comprised of the  $\alpha$ -cardiac myosin heavy chain (MHC) promoter and a modified  $\beta$  galactosidase (nLAC) reporter. To generate the MHC-nLAC transgenic mice, MHC-nLAC insert DNA (see Figure 1) was purified by absorption onto glass beads, dissolved at a concentration of 5  $\mu$ g/ml, and microinjected into the nuclei of one cell inbred C3H3B/FeJ embryos according to established protocols (17). Polymerase Chain Reaction (PCR) analysis was employed to identify founder animals and to monitor transgene segregation. The sense strand primer 5'-GGTGGGGCTCTTACCCCCAGACCTCTCC-3' was localized to the MHC promoter and the antisense strand primer 5'-GCCAGGGTTTTCCAGTCACGACGTTGT-3' was localized to the

nLAC reporter. PCR analyses were as described in (18). The MHC promoter consisted of 4.5 kb of 5' flanking sequence and 1 kb of the gene encompassing exons 1 through 3 up to but not including the initiation codon. The nLAC reporter was modified so as to carry both a eukaryotic translation initiation site and the SV40 nuclear localization signal (19). The mPl sequences carried an intron, as well as transcriptional termination and polyadenylation signals from the mouse protamine 1 gene.

For preparations for examination of  $\beta$  galactosidase ( $\beta$ GAL) activity and DAPI epifluorescence, transgenic animals were heparinized (10,000 U/kg IP) prior to sacrifice by cervical dislocation. Hearts were placed in a beaker of gassed (95% O<sub>2</sub>, 5% CO<sub>2</sub>) KHB buffer (105 mM NaCl, 20 mM NaHCO<sub>3</sub>, 3.8 mM KCl, 1 mM KH<sub>2</sub>PO<sub>4</sub>, 1.2 mM MgSO<sub>4</sub>, 0.01 mM CaCl<sub>2</sub>, 1 mM mannitol, 10 mM taurine, 10 mM dextrose, 5 mM Na-pyruvate). Hearts were then hung by the aorta and perfused with gassed KHB (0.5 ml/min at 37°C) containing 2.5 mM EGTA for five minutes, followed by 0.17% collagenase (Type I, Worthington Biochemical, Freehold NJ) in KHB. Hearts were perfused until flaccid and the ventricles were minced with scissors and isolated cells obtained by triturating with a Pasteur pipette. After at least one hour of formalin fixation, suspensions were filtered and smeared onto positively charged slides (Superfrost Plus, Fisher, Pittsburgh PA), and allowed to dry.

For isolation of single cells for injection, females with 15 day embryos (onset of pregnancy determined by vaginal plugs) were sacrificed by cervical dislocation. Embryos were removed, decapitated, and hearts were harvested under PBS, and ventricles and atria were separated. Transgenic ventricles (identified by cardiac  $\beta$ GAL activity) were digested in 0.1% collagenase

Immediately after isolation, embryonic cardiomyocytes were washed three times with DPBS and directly injected into the ventricular myocardium of syngeneic mice (Jackson Laboratories) under open heart surgery as in Example 1.  $1-10 \times 10^4$  cells were injected in a volume of 2-3  $\mu$ l using a plastic syringe fitted with a 30 gauge needle.

Histology. For H and E, X-GAL, immunohistology and <sup>3</sup>H-thymidine analyses, hearts were removed following cervical dislocation and cryoprotected in 30% sucrose, embedded and sectioned at 10 μm with a cryomicrotome as in Example 1. H and E staining, monitoring for intra-cardiac graft rejection, and assay for [<sup>3</sup>H]-thymidine incorporation were also conducted as in Example 1. To assay βGAL activity, sections were hydrated in PBS, post-fixed in acetone:methanol (1:1) and then overlaid with mixture containing 1 mg/ml X-GAL (5-bromo-4-chloro-3-indolyl-β-D-galactoside), 5 mM potassium ferricyanide, 5 mM potassium ferrocyanide and 2 mM magnesium chloride in PBS. Positive staining is indicated by the appearance of a blue chromophore. After treatment with primary antibody, signal was visualized by an avidin-biotin (ABC) kit (Vector Labs, Burlingame CA). The heart was processed as described above, and sections were post-fixed in methanol:acetone (1:1), stained with H and E, and coated with a thin layer of photographic emulsion (Ilford L4, Polysciences) diluted 1:1 with distilled water. Sections were exposed, developed, washed, fixed and washed as in Example 1. X-GAL staining of single cell preparations was as described above. For visualization of nuclei in single cell preparations,

slides were stained with DAPI in PBS (0.28  $\mu$ M, three min. at room temperature, Boehringer Mannheim), washed three times in PBS, and wet-mounted in 2% propyl gallate dissolved in glycerol. To obtain coronal heart sections, mice were sacrificed by cervical dislocation, hearts were harvested and perfused on a Langendorff apparatus with 2% glutaraldehyde in 0.1 M cacodylate buffer (pH 7.4). After immersion fixation overnight in the same buffer, 200  $\mu$ m coronal sections were made with a vibratome (Campden, London, United Kingdom). To localize the graft, sections were pooled and stained for  $\beta$ GAL activity with X-GAL as described above.

Electron Microscopy. MHC-nLAC embryonic grafts were localized in coronal heart sections as described above. After trimming, the tissue was post-fixed in 2% osmium tetroxide (Stevens Metallurgical Corp., New York NY). Tissue was then dehydrated and embedded in Ladd LX-112 (Ladd Research Industries). Grafted areas were further trimmed, thin sectioned, and stained with uranyl acetate and lead citrate. Specimens were viewed on a Phillips 400 transmission electron microscope as in Example 1.

Electrocardiogram Analyses. ECG analyses were conducted as in Example 1.

## B. RESULTS

Transgenic mice generated as above carried a fusion gene comprised of the MHC promoter and a nLAC reporter. nLAC carries the SV40 nuclear transport signal, which results in the accumulation of  $\beta$  galactosidase activity in the nucleus of targeted cells. Four transgenic lineages were produced, and two (designated MHC-nLAC-2 and MHC-nLAC-4) were selected for further analyses. To ensure

that the MHC-nLAC transgene would provide a suitable cell lineage marker,  $\beta$  galactosidase ( $\beta$ GAL) activity was assessed in transgenic cardiomyocytes. Single cell preparations generated by retrograde collagenase perfusion were examined simultaneously for  $\beta$ GAL activity and DAPI epifluorescence.  $99.0 \pm 0.45\%$  ( $n=400$ ) of the transgenic cardiomyocyte nuclei expressed  $\beta$ GAL, whereas no  $\beta$ GAL activity was detected in noncardiomyocytes. In addition, no nuclear  $\beta$ GAL activity was detected in nontransgenic control cardiomyocytes.

Single cell suspensions were prepared by collagenase digestion of hearts harvested from embryonic day 15 transgenic mice. Greater than 95% of the cardiomyocytes isolated by this technique were viable as evidenced by dye exclusion assay. Cardiomyocytes were delivered to left ventricular free wall of syngeneic nontransgenic animals. Grafted cardiomyocytes were readily and unambiguously identified by virtue of the nuclear  $\beta$ GAL activity encoded by the MHC-nLAC transgene. Grafted cardiomyocytes were frequently observed at sites distal to the point of delivery; it presently is not clear if this distribution of grafted cells reflects cardiomyocyte migration or passive diffusion along dissection planes produced by the injection process. Approximately 50% (7/13) of the animals receiving intra-cardiac injections of embryonic cardiomyocytes developed grafts. This frequency of successful graft formation is likely to increase as cell preparation and implantation protocols are optimized.

Light microscopic analyses of H and E stained sections processed for  $\beta$ GAL activity indicated that grafted cardiomyocytes (blue nuclei) were juxtaposed directly with host cardiomyocytes (purple nuclei). Additional H and E analyses failed to detect significant graft encapsulation. The observed proximity of graft and



host cardiomyocytes and absence of encapsulation are prerequisites for successful coupling between the two cell types.

Consecutive sections of a 19 day old intra-cardiac graft were processed for H and E, BGAL activity, and macrophage and leukocyte immunoreactivity. No evidence for graft rejection was observed, despite the fact that the animals were not immune suppressed. As a positive control for the immunohistology, grafts of incompatible MHC haplotype were produced; graft rejection was clearly evident in these hearts. Tritiated thymidine uptake analyses indicated that only 0.6% (n=156) of the BGAL-positive nuclei were synthesizing DNA, although noncardiomyocyte DNA synthesis was apparent. Since the embryonic day 15 donor cells were still mitotically active when grafted (labeling index of ca. 29%), the exceedingly low level of DNA synthesis observed in BGAL positive cells at 19 days post-grafting suggested that the MHC-nLAC embryonic cardiomyocytes had undergone terminal differentiation.

The juxtaposition of graft and host cardiomyocytes observed by light microscopic analyses prompted a determination whether direct intercellular coupling could be detected between the two cell types. The X-GAL reaction product is an electron-dense precipitate which can be detected by transmission electron microscopy (TEM, see 19). Vibratome sections from glutaraldehyde perfusion-fixed hearts were stained for BGAL activity, and grafted regions thus identified were trimmed and embedded for TEM. BGAL positive nuclei were readily observed by light microscopic analysis of 1  $\mu$ m sections. The X-GAL reaction product had a perinuclear appearance due to a slight degree of nuclear leaching which occurred during the embedding process. The same groups of cardiomyocytes

In addition to documenting the presence of abundant intercellular coupling between grafted and host cardiomyocytes, the TEM analyses revealed that the grafted cardiomyocytes were highly differentiated. Normal characteristics of adult cardiomyocytes were observed including myofibrillae forming complete sarcomeres, numerous junctional complexes between cells and abundant mitochondria. Indeed, aside from the presence of the X-GAL reaction product, grafted cardiomyocytes were indistinguishable from host cells. Further, binucleated, BGAL positive cells could be detected in the intra-cardiac grafts. Because binucleation is a characteristic of adult rodent cardiomyocytes, this observation further supports that the grafted cardiomyocytes have undergone terminal differentiation.

Surface ECG recordings were employed to determine if the presence of coupled embryonic cardiomyocyte grafts negatively influenced host heart automaticity. ECG traces from graft-bearing animals were indistinguishable from sham operated controls, and exhibited P and QRS complexes

typical for mice. There was no evidence for cardiac arrhythmia in graft-bearing animals, despite the presence of a high degree of intercellular coupling between grafted and host cardiomyocytes.

#### EXAMPLE 4

##### Preparation of Substantially Pure Cardiomyocyte Culture

Embryonic stem cells were genetically modified in a manner enabling the production of a substantially homogeneous population of non-immortalized cardiomyocytes. The parental ES cell line (D3) was cotransfected with a pGK-HYG (hygromycin) plasmid and a plasmid containing a MHC-neo<sup>r</sup> gene. The pGK-HYG plasmid provides selection for transfected ES cells, while the mMHC-neo<sup>r</sup> gene facilitates a second round of selection on differentiated cells: incubation in the presence of G418 eliminates non-cardiomyocytes (that is, cells in which the MHC promoter is not active).

Stably transfected ES cells were selected by growth in the presence of hygromycin. The plasmids were linearized and introduced into the stem cells via electroporation at 1180  $\mu$ Farad, 220 volts. The transfected cells were maintained in DMEM supplemented with 10% preselected FBS, 0.1 mM  $\beta$ -mercaptoethanol, nonessential amino acids, PenStrep and LIF, and transformants selected by the addition of hygromycin into the medium. Co-transfectants were then identified by PCR analysis specific for both transgenes. The transfections produced a cell line, designated 9A, which carries both transgenes.

Cardiogenesis was induced in 9A ES cells by plating  $2 \times 10^6$  cells onto uncoated 100 mm bacterial petri dishes in the absence of LIF. After 8 days in culture, numerous patches of cells exhibiting spontaneous contractile

activity (i.e. cardiomyocytes) were observed. At this point, G418 was added to the media, and the cells incubated for an additional 9 days. During this treatment it was apparent that many of the non-cardiomyocytes were being killed by the G418. Importantly, the G418 had no discernible effects on the cardiomyocytes, which retained their spontaneous beating activity throughout the course of the experiment.

After a total of 9 days of G418 selection, the surviving cells were dissociated with collagenase and trypsin, and then replated onto fibronectin coated microscope slides. The cells were cultured an additional 24 hours to allow them to recover from passage, and then fixed for immunocytologic analysis. Cells were reacted with MF20, a monoclonal antibody which recognizes sarcomeric myosin. Cells were then individually counted for the presence or absence of sarcomeric myosin, a marker for cardiomyocytes. The results were as follows:

Total number of cells counted:	794
Number of MF20+ cells:	791
Number of MF20- cells:	3*
Percent cardiomyocytes:	99.6%

\*The 3 cells which did not stain with MF20 may still have been cardiomyocytes, since this antibody will not react with monomeric myosin.

It was thus demonstrated that the expression of drug (neomycin) resistance under a cardiac-specific promoter, enables the selection of an essentially pure population of ES derived cardiomyocytes in culture. Such populations can be used to form myocardial grafts using procedures as discussed in the Examples above.

EXAMPLE 5  
Delivery of Protein via Graft

A. METHODS

C2C12 Cell Culture and transfection. C2C12 myoblasts (ATCC) were maintained in the undifferentiated state by culturing at low density in high glucose Dulbecco's Modified Eagle Media (DMEM)<sup>1</sup> supplemented with 20% fetal bovine serum, 1% chicken embryo extract, 100 units/ml penicillin and 100µg/ml streptomycin. For some studies, myogenic differentiation was induced by culturing in DMEM supplemented with 2% horse serum and antibiotics.

A fusion gene comprised of the metallothionein (MT) promoter driving a modified Transforming Growth Factor-Beta 1 (TGF-β1) cDNA was obtained from Samuel and colleagues (20). Transcriptional activity of the metallothionein promoter can be regulated by modulating the heavy metal content of cell culture media. The TGF-β1 cDNA carried site-directed mutations which resulted in the conversion of Cys<sup>223</sup> and Cys<sup>225</sup> to serines. This modification (described further in (21)) results in the elaboration of a TGF-β1 molecule which is unable to form dimers, and consequently is not subject to normal post translational regulation. Cells expressing the modified cDNA constitutively secrete processed, active TGF-β1 (20). The MT-TGF fusion gene was introduced into C2C12 myoblasts by calcium phosphate transfection; stable transfectants were selected by virtue of co-transfection with an SV40-neo<sup>r</sup> transgene. Four independent clones were isolated, and presence of the transgene was confirmed by Southern blot analysis. The relative levels of TGF-β1 expression in the different clonal cell lines was initially assessed by Northern blot analysis, and one line, designated C2(280), was utilized for subsequent experiments.

000000-000000

Myocardial Grafting Protocol. The grafting protocol was as described in Example 1. Fourteen days post-surgery, graft bearing animals were given heavy metal (25 mM ZnSO<sub>4</sub> in drinking water). Zinc treatment was continued until the termination of the experiment (1-4 weeks).

Histology. For paraffin sections, hearts were fixed in 10% neutral buffered formalin, dehydrated through graded alcohols, and infiltrated with paraffin. Tissue blocks were then sectioned at 6µm. H and E staining was performed directly after sectioning according to manufacturer's specifications (Sigma Diagnostics).

For [<sup>3</sup>H]-thymidine incorporation, mice were given a bolus and sacrificed as in Example 1. The heart was removed and processed for paraffin embedding as described above. Autoradiography was likewise conducted as in Example 1.

## B. RESULTS

Expression of recombinant TGF-β1 in response to heavy metal induction was examined in C2(280) myoblasts and myotubes. Transgene transcripts (1.8 kb) were readily distinguished from those originating from the endogenous TGF-β1 gene (2.5 kb) by Northern blot analysis. Addition of heavy metal to the culture media resulted in a marked increase of recombinant TGF-β1 transcripts in C2(280) myoblasts and myotubes. As indicated above, modified TGF-β1 expressed by C2(280) cells should have constitutive biological activity. To directly test this, conditioned media from C2(280) myoblasts and myotubes was examined by growth inhibition assay.

C2(280) myoblasts were used to produce intra-cardiac grafts in syngeneic C3Heb/FeJ mice. The presence of grafts was readily detected in H and E stained sections. 100% (n > 50) of the animals receiving intra-cardiac injections of C2(280) cells went on to develop grafts. Interestingly, H and E analysis suggested that the C2(280) grafts were somewhat less differentiated as compared to those produced with unmodified C2C12 cells. This result was confirmed by immunohistologic analysis with a monoclonal antibody which recognizes skeletal myosin heavy chain.

C2(280) graft transgene expression was assessed by immunohistology with an anti-TGF- $\beta$ 1 antibody; TGF- $\beta$ 1 expression was readily detected in C2(280) grafts. As a negative control, TGF- $\beta$ 1 expression was assessed in grafts produced by C2C12 myoblasts. As expected, the relative levels of TGF- $\beta$ 1 expression were markedly reduced in C2C12 grafts as compared to C2(280) grafts.

TGF- $\beta$ 1 is a well known angiogenic factor.

<sup>3</sup>H-thymidine incorporation analyses in vascular endothelial cells was therefore assessed to determine if an enhanced angiogenic response occurred in grafts expressing the MT-TGF transgene. DNA synthesis in vascular endothelial cells was readily apparent in C2(280) grafts under administration of a single bolus injection of <sup>3</sup>H-thymidine (H-THY). In contrast, vascular endothelial DNA synthesis was markedly reduced in non-transfected C2C12 grafts (Table 1). To rule out the possibility that the angiogenic responses was due solely to graft mass, <sup>3</sup>H-thymidine incorporation was compared between similar size and aged C2C12 and C2(280) grafts (Table 1). A marked increase in the number of vascular endothelial cells synthesizing DNA was apparent in all of the

09673014-060804  
T06090-T06280

analyses. Finally, the thymidine incorporation assay also revealed that a percentage of the grafted myoblasts continued to proliferate. This observation is consistent with the known inhibitory effect of TGF- $\beta$ 1 on myodifferentiation, and most likely accounts for the undifferentiated appearance of the C2(280) grafts.



TABLE 1  
TGF- $\beta$ 1 Delivery and  
Vascular Endothelial DNA Synthesis

Time Post Zn <u>Induction</u>		C2C12 <u>TGF-<math>\beta</math>1(-)</u>	C2(280) <u>TGF-<math>\beta</math>1(+)</u>
1 Week	Total # $^3\text{H}$ -THY+ Endothelial Cells	0	8
	Total # of Vessel Sections Counted	35	17
	# Synthetic Cells/ Vessel Section	$0.0 \pm 0.00$	$0.46 \pm 0.036$
2 Weeks	Total # $^3\text{H}$ -THY+ Endothelial Cells	1	7
	Total # of Vessel Sections Counted	18	21
	# Synthetic Cells/ Vessel Section	$0.06 \pm 0.056$	$0.34 \pm 0.052$

While the invention has been illustrated and described in detail in the foregoing description, the same is to be considered as illustrative and not restrictive in character, it being understood that only the preferred embodiments have been described and that all modifications that come within the spirit of the invention are desired to be protected.

003304-0600  
T00000-T00000

## REFERENCES

The following references, to the extent that they provide exemplary procedural or other details supplementary to those set forth herein, are specifically incorporated herein by reference.

1. Tompson, L. Fetal transplants show promise. *Science* 257: 868-870, 1992.
2. Gussoni, E., Pavlath, G.K., Lanctot, A.M., Sharma, K.R., Miller, R.G., Steinman, L. and Blau, H.M. Normal dystrophin transcripts detected in Duchenne muscular dystrophy patients after myoblast transplantation. *Nature* 356: 435-438, 1992.
3. Steinhelper, M.E., Lanson, N., Dresdner, K., Delcarpio, J.B., Wit, A., Claycomb, W.C. and Field, L.J. Proliferation in vivo and in culture of differentiated adult atrial cardiomyocytes from transgenic mice. *American Journal of Physiology* 259 (Heart and Circulatory Physiology 28): H1826-H1834, 1990.
4. Janse, M.J., Cinca, J., Morena, H., Fiolet, J.W., Kleber, A.G., deVries, G.P., Becker, A.E. and Durrer, D. The border zone in myocardial ischemia. An electrophysiological, metabolic and histochemical correlation in the pig heart. *Circulation Research* 44:576-588, 1979.
5. Spear, J. F., Michelson, E.L., and More, E.N. Cellular electrophysiologic characteristics of chronically infarcted myocardium in dogs susceptible to sustained ventricular tachyarrhythmias. *Journal of the American College of Cardiology* 4:1099-1110, 1983.

6. Delcarpio, J.B., Lanson, N.A. Jr., Field, L.J. and Claycomb, W.C. Morphological characterization of cardiomyocytes isolated from a transplantable cardiac tumor derived from transgenic mouse atria (AT-1 cells). Circulation Research 69:1591- 1600, 1991.

7. Rockman, H.A., Ross, R.A., Harris, A.N., Knowlton, K.U., Steinhelper, M.E., Field, L.J., Ross, J. Jr. and Chien, K.R. Segregation of atrial-specific and inducible expression of an ANF transgene in an in vivo murine model of cardiac hypertrophy. Proc. Natl. Acad. Sci. USA 88:8277-8281, 1991.

8. Bullock, G.R. and Petrusz, P., in Techniques in Immunocytochemistry, Vol. II, Academic Press, New York, 1983.

9. Field, L.J. Atrial natriuretic factor-SV40 T antigen transgenes produce tumors and cardiac arrhythmias in mice. Science 239:1029-1033, 1988.

10. Katz, E., Steinhelper, M.E., Daud, A. Delcarpio, J.B., Claycomb, W.C. and Field, L.J. Ventricular cardiomyocyte proliferation in transgenic mice expressing  $\alpha$ -Cardiac Myosin Heavy Chain-SV40 T antigen fusion genes. American Journal of Physiology 262 (Heart and Circulatory Physiology 31):H1867-1876, 1992.

11. Steinhelper, M.E. and Field, L.J. "SV40 large T-Antigen induces myocardiocyte proliferation in transgenic mice", in The development and regenerative potential of cardiac muscle, John Oberpriller and Jean Oberpriller, eds. Harwood Academic press, 1990.

FOUO-1103250

12. Steinhelper, M.E. and Field, L.J. Cardiac Tumors and dysrhythmias in transgenic mice. Toxicologic Pathology 18: 464-469, 1990.
13. Yaffe, D., and O. Saxel. Serial passaging and differentiation of myogenic cells isolated from dystrophic mouse muscle. Nature 270:725-727, 1977.
14. Nadal-Ginard, B. Commitment, fusion, and biochemical differentiation of a myogenic cell line in the absence of DNA synthesis. Cell 15:885-864, 1978.
15. Bruni, C. Mitotic activity of muscle satellite cells during the early stages of rhabdomyosarcomas induction with nickel subsulfide. In Muscle Regeneration, A. Mauro, editor. Raven Press, New York, NY, 1979.
16. Rubin, L.L., C.E. Keller and S.M. Schuetze. Satellite cells in isolated adult muscle fibers in tissue culture. In Muscle Regeneration, A. Mauro, editor. Raven Press, New York, NY, 1979.
17. B. Hogan, F. Costantini and E. Lacy, in Manipulating the Mouse Embryo - A Laboratory Manual, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York, 1986.
18. M.E. Steinhelper, K.L. Cochrane and L.J. Field. Hypotension in transgenic mice expressing atrial natriuretic factor fusion genes. Hypertension 16:301-307, 1990.
19. A.D. Loewy, F.C. Bridgman and T.C. Mettenleiter. Brain Research 555:346, 1991.

20. S.K. Samuel, et al. Autocrine induction of tumor protease production and invasion by a metallothionein-regulated TGF-beta 1 (Ser223,225). Embo Journal [JC:emb] 11(4):1599-1605, 1992.

21. A.M. Brunner, et al. Site-directed mutagenesis of cysteine residues in the pro region of the transforming growth factor beta 1 precursor. Expression and characterization of mutant proteins. Journal of Biological Chemistry 264(23):13660-13664, 1989.

22. E.J. Robertson. Embryo-derived stem cell lines, in Teratocarcinomas and embryonic stem cells: a practical approach, E.J. Robertson, editor. IRL Press, Washington DC, 1987.

009704-0000  
FBI/DOJ